

Description

Mazuri® Waterfowl Maintenance Diet has been formulated to meet the nutritional needs of the vast majority of non-specialist feeding ornamental waterfowl.

Features and Benefits

- **Contains cereal grain** – Provides plant-based energy sources.
- **Proper protein level** – Designed to maintain health and vigor.
- **Balanced nutrition/energy ratio** – Aims to help maintain condition.
- **Nutritionally complete** - Requires no mixing or supplementation.
- **Extruded floating pellet diet** - Allows for feeding of bird in the water and prevents accumulation on the bottom of the feeding area, limiting water contamination and waste and creating a more natural feeding environment.
- **Sustainable fishmeal source used.**

Product Form

Extruded pellet size: 3/8" diameter x 3/16" length.

- 50 lb. net weight paper sack

Catalog #

0001435

Guaranteed Analysis

Crude protein not less than	14.0%	Calcium not less than	0.85%
Crude fat not less than	3.0%	Calcium not more than	1.35%
Crude fiber not more than	4.5%	Phosphorus not less than	0.60%
Ash not more than	7.0%	Salt not less than	0.10%
		Salt not more than	0.50%

Ingredients

Ground corn, wheat middlings, ground oats, dehulled soybean meal, calcium carbonate, fish meal (menhaden), soybean oil, dicalcium phosphate, salt, mixed tocopherols (a natural preservative), brewers dried yeast, cholecalciferol (vitamin D₃), menadione sodium bisulfite complex (vitamin K), vitamin A acetate, magnesium oxide, d-alpha tocopheryl acetate (natural source of vitamin E), choline chloride, zinc oxide, calcium pantothenate, niacin, copper sulfate, marigold extract, dl-methionine, vitamin B₁₂ supplement, l-lysine, calcium iodate, riboflavin, folic acid, sodium selenite, biotin.

Feeding Directions

Feed ad libitum to non-breeding mature, fully feathered birds.

The feed can be offered ad libitum to non-breeding mature, fully feathered birds.

Adult birds should eat in the range of 25-30 grams/kg of body weight each day, depending on species size.

Scatter directly on the water, where it will float, or it can be fed in a long, shallow trough, which will prevent the food from floating away from the birds.

If supplementation is desired, provide species-specific food items at not more than 20% of the total dietary intake (as is basis).

Approximate Nutrient Composition¹**NUTRIENTS**

Protein, %	14
Arginine, %.....	0.87
Cystine, %.....	0.28
Glycine, %.....	0.66
Histidine, %.....	0.38
Isoleucine, %.....	0.56
Leucine, %.....	1.2
Lysine, %.....	0.69
Methionine, %.....	0.27
Phenylalanine, %.....	0.64
Tyrosine, %.....	0.38
Threonine, %.....	0.50
Tryptophan, %.....	0.15
Valine, %.....	0.67

Fat (Acid hydrolysis), %	4.5
Linoleic acid, %.....	1.7

Fiber (Crude), %	4.5
Neutral Detergent Fiber, %.....	18
Acid Detergent Fiber, %.....	5.5

Metabolizable Energy²,	
kcal/kg.....	2,830

MINERALS

Ash, %	5.6
Calcium, %.....	1.1
Phosphorus, %.....	0.60
Phosphorus (non-phytate), %.....	0.29
Potassium, %.....	0.64
Magnesium, %.....	0.20
Sodium, %.....	0.19
Chloride, %.....	0.29
Iron, ppm.....	175
Zinc, ppm.....	90
Manganese, ppm.....	108
Copper, ppm.....	10
Iodine, ppm.....	1.3
Selenium (added), ppm.....	0.30

VITAMINS

Thiamin, ppm.....	5.9
Riboflavin, ppm.....	2.6
Niacin, ppm.....	38
Pantothenic acid, ppm.....	16
Choline, ppm.....	1,000
Folic acid, ppm.....	1.3
Pyridoxine, ppm.....	4.2
Biotin, ppm.....	0.24
Vitamin B ₁₂ , µg/kg.....	9.3
Vitamin A, IU/kg.....	8,940
Vitamin D ₃ (added), IU/kg.....	2,135
Vitamin E, IU/kg.....	85
Vitamin K (as menadione), ppm.....	2.5

Storage Conditions

For best results, store contents of open bag in container with sealing lids. Store in a cool (75°F or colder), dry (approximately 50% RH) location. Use within 1 year of bag manufacturing.

¹ Based on the latest ingredient analysis information. Since nutrient composition of natural ingredients varies, analyses will vary accordingly.

² Calculated using data from domesticated poultry.

Mazuri® is a registered trademark of Purina Mills, LLC.